

حمل الآن

مجاناً وحصرياً


# المراجعة رقم (1)

## الترم الاول



# physics

- Chapter 1 Wave Motion
- Chapter 2 Light
- Chapter 3 Hydrodynamics



فیزکس ٹیچر  
PHYSICS  
TEACHER





# Types of motion



1



2




## Translational motion



It's the motion that has starting point and ending point

1

motion in straight-line



2

motion in curved path

(projectile motion)



## Periodic motion



It's the motion that repeats itself regularly in equal intervals of time. Periodic motion can be classified

into:

1

Wave motion



2

Oscillatory motion





# Oscillatory motion



1

Vibratory motion

2

Amplitude

3

Displacement

4

Complete vibration

أونلاين







# Oscillatory motion



## 1 Vibratory motion (oscillatory motion)

It's the **motion made by a particle** on both sides away from its rest position.






# Oscillatory motion



2

## Amplitude



It's the **maximum displacement** done by a vibrating body away from its rest position.

Or It's the **distance between to points** where the velocity at one point is maximum & at the other is zero.

Or It's the **distance between the position** at which K.E is maximum & P.E is zero & other position of contra conditions.







# Oscillatory motion



3

## Displacement

It's the distance covered by the body between its position at any instant & its position at rest.







# Oscillatory motion



4

## Complete vibration or (Wave form)

It's the disturbance when a vibrating particle passes by a fixed point two successive times in the same direction.







## Periodic time "T"



It's the time taken by a vibrating body to make one complete vibration or cycle.

$$T = \frac{t \text{ (time in second)}}{n \text{ ( number of complete oscillations)}}$$

$$T = 4 \times \text{Time of amplitude}$$



## Frequency " : ( Hertz or cycle/ sec )



It's the no of complete **cycles or vibrations** done by a vibrating body in one second

Its unit is **K.HZ** =  $10^3$  Hz & **Mega Hz** =  $10^6$  Hz



$$\nu = \frac{n(\text{number of oscillations})}{t(\text{time})}$$







## Phase



It is the **position & velocity and direction** of a certain molecule at a certain instant



Difference:  $0^\circ$




In Phase



## Important notes



If two **bodies vibrate** having the same phase that means that they have the same **velocity & direction** as the following



(a) & (d)

have the same **direction & velocity**

(b) & (c)

have the **same direction** but not the same velocity.





# Example for vibrating bodies



## (simple harmonic motion)


### Motion of a simple pendulum

In which the displacement increases from the starting point (A) to +ve maximum value at (B) then it decreases to zero at (A) then increases to -ve max value at (C) then decreases to zero at (A) and repeat itself through uniform intervals of time.

$$E = KE + P.E$$

$$K.E = \frac{1}{2} mv^2$$

$$P.E = mgh$$






# Oscillatory motion graph



Position	A	B	C
Displacement	0	Y	-Y
Velocity	Max	0	0
P.E	Min	Max	Max
K.E	Max	0	0









## Notes

1

if the body starts motion from the position at which the velocity is zero then the curve is drawn as a sin curve

2

If the body starts its motion from the position at which the velocity is maximum then the curve is a cos curve.






**In the opposite figure ....**

**If the time taken by the pendulum to move from A to C is 0.8 s, calculate**

**1) The periodic time**



☐ 0.6

☐ 2

☐ 1.6

☐ 5

هنذاكر  
أونلاين






**In the opposite figure ....**

**If the time taken by the pendulum to move from A to C is 0.8 s, calculate**

**2) The frequency**



☐ 0.625

☐ 2

☐ 0.6

☐ 5


هنذاكر  
أونلاين



**In the opposite figure ....**

**If the time taken by the pendulum to move from A to C is 0.8 s, calculate**

**3) The number of complete oscillations through 16s.**



☐ 15

☐ 10

☐ 9

☐ 13

فنذاکر  
اونلاين






**In the opposite figure ....**

**If the time taken by the pendulum to move from A to C is 0.8 s, calculate**

**4) The time required to make 50 oscillations**



☐ 80s

☐ 60s

☐ 75 s

☐ 60s

ہندذاکر  
آن لائن



يعنى ايه موجة ؟

## Wave

It's a **disturbance that can propagate transferring energy** in the direction of propagation without any transfer in the medium particles.







## Mechanical Waves & Electromagnetic waves



Mechanical Waves	Electromagnetic waves
It's a <b>disturbance</b> needs a <b>medium</b> to propagate.	It's the <b>disturbance</b> propagates in <b>space and other media</b> .
It's formed from <b>transverse</b> or <b>longitudinal</b> waves.	It's formed from <b>transverse waves only</b> .
Formed from the vibration of the particles of the medium.	Formed from the vibration of electric & magnetic fields, which is perpendicular to each other & perpendicular to the direction of wave propagation.




# Mechanical waves




1

Water waves




2

waves produced through  
vibrating strings



3

sound waves








# Electromagnetic waves




1

Light waves



2

Y rays



4

X rays

3

Radio waves





## Mechanical wave





It is a **disturbance that propagates** in materialistic media



## Electromagnetic waves



They are **waves originated** from vibrating electric and magnetic fields having the same phase with frequency ( $\nu$ ), perpendicular to each other and to the direction of the wave propagation and can spread in materialistic media and in space.







## Give reasons



1

**We see the light produced from the cosmic explosions and we don't hear the sound produced from them.**

**Because**

**the light is an electromagnetic wave that can propagate through the space while the sound is a mechanical wave that needs a medium to propagate and cannot propagate through space.**



**Give reasons**



**2**

**Astronauts use wireless devices on the moon surface.**

**Because**

**there is no medium on the moon, so the sound cannot propagate through space.**

**Or**

**sound is a mechanical wave that needs medium to propagate and cannot propagate through space.**





**Give reasons**



**3**

**The electromagnetic waves propagate through space.**

**Because**


**it consists of two fields electric and magnetic perpendicular to each other where both of them don't need medium to propagate through.**



## Conditions required for producing mechanical waves



- 1 Vibrating source (**pendulum – Fork - stretched wire**).



- 2 A disturbance to transfer energy from the source **to the medium**.
- 3 A medium that carry the **disturbance**.





# Mechanical waves



1



Transverse waves




Transverse Wave



2

Longitudinal waves



Longitudinal Wave



اولاين



# Transverse waves & Longitudinal waves

## 1 Transverse waves

## 2 Longitudinal waves

<b>Wave form</b>	 <p>A diagram of a transverse wave represented by a red sine wave. The wave oscillates vertically. Two peaks are labeled 'Crest' and two valleys are labeled 'Trough'. The horizontal distance between two consecutive crests is marked with a double-headed arrow and labeled <math>\lambda</math>. Similarly, the horizontal distance between two consecutive troughs is also marked with a double-headed arrow and labeled <math>\lambda</math>.</p>	 <p>A diagram of a longitudinal wave represented by vertical purple lines of varying density. The wave oscillates horizontally. Regions with fewer lines are labeled 'Rarefaction' and regions with more lines are labeled 'Compression'. The horizontal distance between the centers of two consecutive rarefactions is marked with a double-headed arrow and labeled <math>\lambda</math>. Similarly, the horizontal distance between the centers of two consecutive compressions is also marked with a double-headed arrow and labeled <math>\lambda</math>.</p>
<b>Direction of vibration of medium particles</b>	Perpendicular to the direction of wave propagation.	Along the line of wave propagation.
<b>Composition</b>	Crests and troughs	Compressions and rarefactions
<b>Wavelength</b>	The distance between two successive crests or two successive troughs.	The distance between the centers of two successive compressions or the centers of two successive rarefactions.
<b>Examples</b>	<ul style="list-style-type: none"><li>• Propagating waves in strings.</li><li>• Waves on water surface.</li></ul>	<ul style="list-style-type: none"><li>• Sound waves in gases.</li><li>• Waves inside water.</li></ul>



$$T = \frac{1}{v} = \frac{\text{Time in second (t)}}{\text{Number of complete vibrations (n)}}$$

$T = 4 \times \text{Time of the amplitude}$

**Periodic time**

**Frequency**

**Wave length**

**waves**

$$\lambda = \frac{\text{Total distance (x)}}{\text{Number of waves (n)}}$$

$$v = \frac{\text{Number of waves (n)}}{\text{Time in seconds (t)}}$$

$$V = \lambda v = \frac{\lambda}{T}$$

**If the two waves have the same**

**Velocity**

**(Same medium)**

$$\frac{\lambda_1}{\lambda_2} = \frac{v_2}{v_1}$$



$$\frac{V_1}{V_2} = \frac{\lambda_1}{\lambda_2}$$

**Frequency**

**(Two different media)**



1

## Main Point

The propagation of the transverse wave in a medium requires the ability of molecules of the medium to vibrate and the existence of a certain cohesive forces between the molecules of the medium, therefore it is transfer in liquid and in solids, but not propagate in the gases.

ہندذاکر  
اؤنلاین





## Main Point

2

When wave is occurs **in water**, then:

on the surface, there is a transverse wave .

**a** (Due to being of cohesive forces between the molecules of water, they vibrate perpendicular to direction of wave propagation to form crests and troughs).

At the depth, there is a longitudinal wave.

**b** (Because the attraction force between the molecules of water at the bottom is minimum value or weak , so they vibrate along the direction of wave propagation to form compressions and rarefactions).

## Main Point

3

# To calculate wavelength

$$\lambda = \frac{x}{N}$$

$x$  : displacement covered by wave.  
 $N$  : number of waves





4

## Main Point

Calculation of **number of wavelengths**, starting from the crest or the trough to another one of the same phase then:

$$\text{no of wavelengths} = \text{no greater order} - \text{no lowest order}$$



5

## Main Point

Calculation of **number of wavelengths**, starting from the crest to the trough then:

**no of wavelengths=**

**(no greatest order - no lowest order) + 1/2**





6

## Main Point

Calculation of number of wave lengths, starting from the trough to the crest then:

$$\text{no of wavelengths} = (\text{no greatest order} - \text{no lowest order}) - 1/2$$



## Main Point

7

# The number of waves (**N**)

$$N = \frac{x}{\lambda} = \frac{t}{T}$$





8

## Main Point

**Amplitude of the standing wave**  
**= the vertical distance between crest and trough.**

هنذاكر  
أونلاين



1

The figure shows a transverse wave, Find

The wave **length** -- The **period time** -- The **frequency**

-- The **amplitude** -- The **velocity**


1 Length = .....

2 period time = .....

3 frequency = .....

4 amplitude = .....

5 velocity = .....







2

Two tones have frequencies **680 Hz and 425 Hz** propagate in the same medium. If the wavelength of one of them exceeds the wavelength of the other wave **by 30 cm**, calculate the velocity of sound in air

ہندذاکر  
اؤنلاین



A girl stood on the beach to watch the waves. She observed that every two seconds, four waves pass in front of her and each wave has a length of **0.5 m**. So, the wave velocity is .....


- ☐ 0.2 m/s
- ☐ 0.25 m/s
- ☐ 0.5 m/s
- ☐ 1 m/s

ہندذاکر  
آن لائن





If the opposite graph shows the relation between frequency and wavelength for a range of frequencies different .....of ultrasonic waves through two media (A , B) , so



☐  $V_A > V_B$

☐  $V_A < V_B$

☐  $V_A = V_B$

☐ not enough data for conclusions



If the time interval between the first crest and the tenth crest in a wave motion is **0.2 s**, then the frequency is.

☐ 45Hz

☐ 50Hz

☐ 55Hz

☐ 60Hz

ہندذاکر  
آن لائن





**In the opposite figure:**

**Points ..... Have the same phase.**



☐ (a,b,c)

☐ (b,c)

☐ (a,b)

☐ (b,d)



Two sound waves of frequencies **300 Hz**, **600 Hz** travels in air, so the ratio between their speeds is .....

☐  $\frac{1}{2}$

☐ 1

☐ 2

☐  $\frac{1}{4}$

ہندذاکر  
اُونلاين






A drop of water fell on the surface of a still water, so a **120** wave ripples are produced during half minute. If the radius of the external circle is **3 m**, the speed of propagation of the wave equals .....

- ☐ 0.01 m/s
- ☐ 0.1 m/s
- ☐ 0.5 m/s
- ☐ 1 m/s

هنذاكر  
أونلاين



The opposite (**displacement – time**) graph depicts the motion of one point in a transverse wave transmission medium, so .....



☐ A

☐ B

☐ C


☐ D

Amplitude (A) cm	Frequency ( $\nu$ ) Hz
6	2.5
6	0.4
3	1.25
3	0.8





The opposite figure shows a wave which is traveling through a medium (x) then it moves to another medium (y), so the ratio between the speed of the wave in medium (x) to its speed in medium (y) is .....



- ☐ greater than one
- ☐ less than one
- ☐ one
- ☐ indeterminable



A wave of frequency  $\nu_1$  and wavelength  $\lambda_1$  propagates in a medium with speed  $v_1$  if this wave travels from this medium to another medium where its speed becomes  $\frac{2}{3} v_1$ , then ....

- the frequency  $\nu_1$  remains constant and the wavelength becomes  $\frac{3}{2} \lambda_1$ .
- the frequency  $\nu_1$  remains constant and wavelength becomes  $\frac{2}{3} \lambda_1$ .
- the wavelength  $\lambda_1$  remains constant and the frequency becomes  $\frac{3}{2} \nu_1$
- the wavelength  $\lambda_1$  remains constant and the frequency becomes  $\frac{2}{3} \nu_1$





**Disturbance that propagates and transfers the energy  
in the same direction of wave the propagation**

- **wave**
- **mechanical wave**
- **complete oscillation**
- **displacement**

A photograph of a bright sun shining through a dense canopy of trees. The sun is positioned in the center, creating a strong lens flare effect that radiates across the frame. The sky is filled with large, white, fluffy clouds. The trees in the foreground are dark and silhouetted against the bright light of the sun. The overall scene is a natural, outdoor setting, likely a park or a wooded area, captured during the day.

**How do we see the light produced from the cosmic  
Explosion as a sun**



A close-up photograph of a human eye, looking directly at the camera. A vibrant rainbow light effect is superimposed over the eye, with colors ranging from red to violet. The light appears to be emanating from the eye itself, creating a glowing, ethereal effect. The background is a soft, out-of-focus skin tone.


**How can we see an objects ?**



# Electromagnetic waves




They consist of two vibrating fields, electric and magnetic field having same frequency, same amplitude and are perpendicular on each other and to the direction of propagation.







# Electromagnetic spectrum



It is the distribution and arrangement of electromagnetic waves according to their wavelengths or their frequencies.

The visible spectrum from  $4000 \text{ \AA}$  to  $7000 \text{ \AA}$

Infra red (IR) more than  $7000 \text{ \AA}$  and Ultra violet (UV) less than  $4000 \text{ \AA}$ .



**HOW**  
**Can see ?**  
**an objects**







# Properties of visible light waves



**1** Light **Propagation**


**2** Light **Reflection**

**3** Light **Refraction**

**4** Light **Interference**

**5** Light **Diffraction**

# 1 Light Propagation








## Properties of visible light waves




### 2 Light Reflection



## Tracing

Three plane mirrors are found as three sides of a square as in the opposite figure. Trace the path of a light ray that falls on one of them until its reflection from the three mirrors.





**Trace (Follow) the path of the light ray**





Fig (1)




هنذاكر  
أونلاين



## Properties of visible light waves



### 3 Light Refraction







## Main Point

1

The light refracted between two optical media due to the change in the velocity of light in these media, so that.

$${}_1n_2 = \frac{v_1}{v_2} = \frac{\sin \phi}{\sin \theta} = \frac{\lambda_1}{\lambda_2} = \frac{n_2}{n_1}$$




2

## Main Point

When light ray travels from **less density medium** to **dense medium**, the velocity of it **decrease**, then it refracts towards the normal.

$$V_1 > V_2 \rightarrow \therefore \sin \phi > \sin \theta \rightarrow (\phi > \theta)$$

$$\therefore \lambda_1 > \lambda_2 \quad \& \quad {}_1n_2 > 1$$






## Main Point

3

When light ray travels from **dense medium** to **less density medium**, the velocity of it increase, then it refracts away the normal:

$$V_1 < V_2 \rightarrow \therefore \sin \phi < \sin \theta \rightarrow \phi < \theta$$

$$\therefore \lambda_1 < \lambda_2 \quad \& \quad n_1 < n_2$$






4

## Main Point

If the incidence ray is **normal** to the surface of separation.

$$\phi = 0^\circ \rightarrow \therefore (\sin \theta = \frac{\sin \phi}{1n2} = \frac{\sin \theta}{1n2} = 0) \rightarrow \theta = 0^\circ$$

It will pass in the same direction of incidence, without refraction.







5

## Main Point

The velocity of light in free space (air) is universal constant and equal to  $c = 3 \times 10^8 \text{ m/s}$  which is greater than the velocity of light ( $V$ ) in any other medium. ( $C > V$  usually).


ہندذاکر  
اُونلاين



# Properties of visible light waves




## 4 Light Interference








## 5 Light Diffraction



# Light

## Light reflection




Angle of incidence  
=  
Angle of reflection

## Light interference young's experiment

Distance between two  
fringes of the same kind

$$\Delta y = \frac{\lambda R}{d}$$




## Light refraction

Snell's Law  $n_1 \sin \phi = n_2 \sin \theta$

Absolute refractive index for a medium

$$n = \frac{c}{v} = \frac{\sin i}{\sin r}$$



Relative refractive index between two media

$${}_1n_2 = \frac{1}{{}_2n_1} = \frac{n_2}{n_1} = \frac{V_1}{V_2} = \frac{\sin i}{\sin r}$$





If the opposite table shows some selected wavelenths from the electromagnetic spectrum in air, so .....

☐  $M < Z < Y$

☐  $Y < Z < M$


☐  $Y < Z = M$

☐  $Y = Z > M$

	Wavelength
visible light	M
Gamma rays	Y
X-rays	Z



The opposite figure shows a light ray which is **falling on the surface of a plane mirror** and bouncing back, hence the angle of reflection of the ray from the mirror equals



.....

☐  $40^\circ$

☐  $50^\circ$

☐  $80^\circ$

☐  $100^\circ$

ہندذاکر  
اؤنلاین





When a light ray transmits from a less optical dense medium to another optical denser medium with an angle of **incidence = zero**, which of the following light properties does not change? .....

- ☐ Speed
- ☐ Wavelength
- ☐ Direction
- ☐ No correct answer



When a light ray falls from air on water surface at an angle of  $60^\circ$ , the angle of refraction will be .....

- ☐ greater than  $60^\circ$
- ☐ less than  $60^\circ$
- ☐ equal to  $60^\circ$
- ☐ equal to  $0^\circ$





When a light ray falls on the interface between two media at an angle of  $\frac{\sin \phi}{\sin \theta}$  incidence  $\phi$  and refracts at an angle of refraction  $\theta$ , the ratio is .....

- ☐ constant for the two media
- ☐ variable according to the value of
- ☐ constant and greater than one
- ☐ constant and less than one







When the **angle of incidence on the boundary** surface between two media doubles, the relative refractive index of them .....

- ☐ decreases to half
- ☐ doubles
- ☐ remains constant
- ☐ increases two times





**Which of the following graphs represents the relation between the relative refractive index between two media and the angle of refraction in one of them?**









**In Young's double –slit experiment, if  $R = 10^4$  d so .....**

- ☐  $\Delta y = \lambda$
- ☐  $\Delta y = 10^4 \lambda$
- ☐  $\Delta y = 10^{-4}$
- ☐  $\Delta y = \lambda/10$





Which of the following **diagrams represents** correctly the phenomenon of light diffraction when light falls on an aperture? .....





**Light waves fall on different aperture of different sizes ,  
so the diffraction of light will be most observable if the  
aperture size is.....**



1m



$10^{-2}\text{m}$



$10^{-3}\text{ m}$



$10^{-5}\text{ m}$

ہندذاکر  
آن لائن





A light ray falls on the separating surface between two media. If the angle between the incident ray and the separating surface is  $40^\circ$  and the angle of refraction in the second medium is  $30^\circ$ . **calculate the relative refractive index from the first medium to the second one.**

ہندذاکر  
اؤنلاین



A light ray falls to water with **angle  $45^\circ$**  . Determine the direction of both the reflected and refracted rays.

(knowing that the refractive index of water is 1.35)

هنذاكر  
أونلاين





In the double-slit experiment; the distance between the two slits is **0.00015m**, the distance between the double-slit screen and the observation screen is **0.75m** and the distance between two successive bright fringes is **0.003m**.


**Calculate the wavelength of the used monochromatic light source.**

هنذاكر  
أونلاين



**From the opposite figure**

**Find the value of each of the refraction and reflection angles.**





هنذاکر  
أونلاين






## The total reflection & the critical angle



اونلاين




## Critical angle between two media



$$\sin \phi_c = \frac{n_{\text{less}}}{n_{\text{more}}} = \frac{\sin(\phi_c)_{\text{more}}}{\sin(\phi_c)_{\text{less}}}$$

In case that the less optically dense medium is air:



$$\sin \phi_c = \frac{1}{n}$$





## Critical angle between two media



هنذاکر  
اونلاين



**An optical fiber that has a material of refractive index 2.1, is coated by an external layer, so the refractive index of the external layer that makes the critical angle between the two layers equal  $32^\circ$  is .....**

☐ 1.11

☐ 1.9

☐ 3.96


☐ 4.32

ہندذاکر  
اُونلاين






The opposite figure shows a light source that is placed inside a transparent medium. so what happens to ray 3 at the boundary surface between the two media?



- ☐ It gets reflected because the angle of incidence is less than the critical angle between the two media
- ☐ It gets reflected because the angle of incidence is greater than the critical angle between the two media
- ☐ It gets refracted because the angle of incidence is less than the critical angle between the two media
- ☐ It gets refracted because the angle of incidence is greater than the critical angle between the two media



The opposite figure shows the occurrence of mirage, hence the correct order of refractive indices of air layers is.....



☐  $n_3 < n_2 < n_1$

☐  $n_1 < n_2 < n_3$

☐  $n_2 < n_1 < n_3$

☐  $n_3 < n_1 < n_2$

ہندذاکر  
اُونلاين






**If you have two flexible transparent materials** where the refractive index of one of them is larger than that of the other material and we want to use them to make an optical fiber, then which of them is used to make the inner core of the optical fiber and which of them is used to make the external layer of it? **And why?**

هنذاكر  
أونلاين



The opposite figure shows four light rays that fall on an isosceles triangular prism of refractive index 1.5, so which of these rays changes its direction **by  $180^\circ$** ?



ہندذاکر  
آن لائن









**The total internal reflection** for an incident light ray from an optically denser medium to an optically less dense medium, occurs when the angle of incidence is .....

- ☐ equal to  $90^\circ$
- ☐ bigger than the critical angle
- ☐ equal to the critical angle
- ☐ less than the critical angle



figure ..... Represent the correct path of incident light ray on a semi-circular disc of glass with an angle of incidence equal the critical angle ( $\phi_c$ )







هنذاكر  
أونلاين

Angle of deviation  $\leftarrow$

**Normal Prism**


$\rightarrow$  Apex angle

$$\alpha = \phi_1 + \theta_2 - A$$

$$A = \theta_1 + \phi_2$$



**If**



**light ray  
falls normally**

$$\phi_1 = \theta_1 = 0^\circ$$

$$A = \phi_2$$

$$\alpha = \theta_2 - \phi_2$$

**Emergent ray  
is tangent**

$$\theta_2 = 90^\circ$$

$$\phi_2 = \phi_c$$

$$A = \theta_1 + \phi_c$$

$$\alpha = \phi_1 + 90 - A$$

**Emergent ray  
is normal**

$$\phi_2 = \theta_2 = 0^\circ$$

$$A = \theta_1$$

$$\alpha = \phi_1 - \theta_1$$



## Position of minimum deviation


Angle of incidence and refraction in the second medium	Refractive index of prism	Angle of deviation	Angles of incidence and refraction in the first medium
$\theta_1 = \phi_2 = \theta_o = \frac{A}{2}$	$n = \frac{\sin \left( \frac{\alpha_o + A}{2} \right)}{\sin \left( \frac{A}{2} \right)}$	$\begin{aligned} \alpha_o &= 2\phi_1 - A \\ &= 2\theta_2 - A \\ &= 2\phi_o - A \end{aligned}$	$\phi_1 = \theta_2 = \phi_o = \frac{\alpha_o + A}{2}$



Light ray falls perpendicular one side of a triangular prism of refractive index **1.5** as shown in figure.

**1** Trace the path of the light ray inside the prism

**2** Find its angle of emergence







A light ray falls on a side of an equilateral triangular prism by angle  $45^\circ$ . Its refractive index is  $\sqrt{2}$ , calculate the angle of emergence and angle of deviation of the ray

ہندذاکر  
آن لائن



The angle of a triangular prism is  $45^\circ$  if a ray of light falls perpendicular on one side then it emerges as tangent to the other side. Calculate the refractive index of the prism material

ہندذاکر  
آن لائن






### The opposite graph:

Represents the relationship between the incidence angle ( $\phi_1$ ) and the deviation angle ( $\alpha$ ) of a light ray falling on the face of a triangular prism. Using the values shown in the figure, Calculate:

1 The emergence angle of the ray at point x .

2 The apex angle of the prism .

3 The refractive index of the prism material .



## Thin Prism

Angle of deviation



$$\alpha_o = A (n - 1)$$

Angular dispersion



$$(\alpha_o)_b - (\alpha_o)_r = A (n_b - n_r)$$

Average deviation



$$(\alpha_o)_y = \frac{(\alpha_o)_b + (\alpha_o)_r}{2} = A (n_y - 1)$$



## Thin Prism

Dispersive power

$$\rightarrow \omega_a = \frac{(\alpha_o)_b + (\alpha_o)_r}{(\alpha_o)_y} = \frac{n_b - n_r}{n_y - 1}$$

Average refractive index

$$\rightarrow n_y = \frac{n_b + n_r}{2}$$




A thin prism of angle  $8^\circ$ , the refractive index of its material to the red and blue colors are 1.52 and 1.54,  
**calculate:**

- 1 The deviation angle for each color
- 2 The angular dispersion between two colors
- 3 The dispersive power of the prism





A light ray falls on a triangular prism and emerges normal to the other face as in the opposite figure, so the angle of incidence ( $\phi$ ) is.....



- ☐ greater than A
- ☐ less than A
- ☐ equal to A
- ☐ there is no relation between it and A



A triangular prism of apex angle  $45^\circ$  and refractive index 1.6 is set on the minimum deviation position, so the angle of incidence of the light ray equals .....

☐  $13.8^\circ$

☐  $17.3^\circ$

☐  $30.5^\circ$

☐  $37.8^\circ$

ہندذاکر  
آن لائن





**The factor (s) that affect the angle of deviation of the light ray in a triangular prism is (are).....**

- ☐ the apex angle of the prism
- ☐ the angle of incidence of the light ray
- ☐ the refractive index of the prism
- ☐ all the previous



A triangular prism of apex angle  $45^\circ$  and refractive index 1.66 is submerged in a liquid of refractive index 1.33. If the prism is in the minimum deviation position. the angle of deviation of light in the prism in this case equals .....


- ☐ 9.29°
- ☐ 12.06°
- ☐ 16.19°
- ☐ 33.88°

ہندذاکر  
آن لائن





The opposite graph represents the relation between the first angle of refraction ( $\theta_1$ ) and the second angle of incidence ( $\phi_2$ ) when a light ray passes through a triangular prism. If the critical angle of the prism material is  $41.8^\circ$ , then the angle of minimum deviation for the falling light ray is .....



☐  $17.27^\circ$

☐  $21.73^\circ$

☐  $25.46^\circ$

☐  $30.25^\circ$

ہندذاکر  
آن لائن



**In the opposite figure, the apex angle of the triangular prism is**


.....

☐  $30^\circ$

☐  $60^\circ$

☐  $70^\circ$

☐  $90^\circ$




هنذاكر  
أونلاين





In the opposite figure shows a triangular prism where a light ray falls on its face at an angle ( $\theta_1$ ) and emerges from the other face at an angle ( $\theta_2$ ), so the ratio equals.....



$$1/2$$



$$1/1$$



$$1/\sqrt{2}$$



$$1/\sqrt{3}$$

ہندذاکر  
آنونلااین

# Laws and solved examples

Volume



Flow rate



Mass

Volume of fluid  
Flowing in time (1)

$$V_{ol} = Q_v t$$

$$Q_v = Av$$

Mass of fluid  
Flowing in time (1)

$$m = Q_m t$$

$$\rho Q_v = \rho Av$$



# Continuity equation

For a tube having two cross – sectional areas

$$A_1 V_1 = A_2 V_2$$

$$r_1^2 V_1 = r_2^2 V_2$$

For a tube branched into a number of branches

Same cross-sectional area

$$A_1 V_1 = \eta A_2 V_2$$

$$r_1^2 V_1 = \eta r_2^2 V_2$$

Different cross-sectional area

$$A_1 V_1 = A_2 V_2 + A_3 V_3 + \dots$$

$$r_1^2 V_1 = r_2^2 V_2 + r_3^2 V_3 + \dots$$



**Three water taps**, the first fills a tank in one hour and the second fills the same tank in half an hour. While the third fills it in quarter an hour. Calculate the time required to fill the tank if the three taps are opened at the same time.

ہندذاکر  
آن لائن






A major artery, the speed of blood in the main artery is  $0.08 \text{ m/s}$  and it branches into **150 blood** capillaries. If the radius of each is  $\frac{1}{8}$  the radius of the main artery. **Calculate the speed of blood in each capillary**

هنذاكر  
أونلاين



viscosity



1



Force of viscosity



2

Viscosity coefficient



فنذا  
أونلاين





A plate of area  $0.01 \text{ m}^2$  moves in a speed of  $12.5 \text{ cm/s}$  parallel to another static plate and isolated from it with a layer of liquid of thickness  $2 \text{ mm}$ , if the viscosity coefficient of the liquid is  $4 \text{ kg/m.s}$ .

Calculate the required force to keep the plate moving

هنذاكر  
أونلاين



In the steady **flow of liquid**, the ratio between the number of streamlines passing in the wide part of a tube to that in the narrow part of the same **tube** is .....

- ☐ greater than one
- ☐ less than one
- ☐ equal to one
- ☐ indeterminable





In the **steady** flow, when the cross-sectional area of the tube decreases, the density of the **streamlines** .....

- ☐ Increases
- ☐ Decreases
- ☐ Vanishes
- ☐ Remains unchanged



If the **cross-sectional area** of tube increases to the double in the steady flow, the **volume flow rate** .....

- ☐ Increases to the double
- ☐ Decreases to the half
- ☐ Remains constant
- ☐ Decreases to the quarter





A water pipe of cross-sectional area  $4\text{cm}^2$  at the ground floor and  $2\text{cm}^2$  at the upper floor. If the water is flowing steadily such that its speed at the ground floor was  $2\text{m/s}$ , then (Where: the water density =  $1000\text{ kg/cm}^3$ )

The flow speed at the upper floor is .....

☐ 1 m/s

☐ 2 m/s

☐ 3 m/s

☐ 4 m/s

ہندذاکر  
اؤنلاین



A water pipe of cross-sectional area  $4\text{cm}^2$  at the ground floor and  $2\text{cm}^2$  at the upper floor. If the water is flowing steadily such that its speed at the ground floor was  $2\text{m/s}$ , then (Where: the water density =  $1000\text{ kg/cm}^3$ )

The volume flow rate of water at the ground floor equals .....

- ☐  $4 \times 10^{-4} \text{ m}^3/\text{s}$
- ☐  $6 \times 10^{-4} \text{ m}^3/\text{s}$
- ☐  $8 \times 10^{-4} \text{ m}^3/\text{s}$
- ☐  $12 \times 10^{-4} \text{ m}^3/\text{s}$





A water pipe of cross-sectional area  $4\text{cm}^2$  at the ground floor and  $2\text{cm}^2$  at the upper floor. If the water is flowing steadily such that its speed at the ground floor was  $2\text{m/s}$ , then (Where: the water density =  $1000\text{ kg/cm}^3$ )

The mass flow where rate of water at the upper floor equals .....

☐ 1.2 kg/s

☐ 0.8 kg/s

☐ 0.6 kg/s

☐ 0.4 kg/s

ہندذاکر  
آن لائن



The ratio between **the mass flow rate** and **the volume flow rate** for a liquid which flows steadily **equals .....**

- The cross-sectional of the tube
- The time of liquid flow
- The speed of liquid flow
- The liquid density









**Water is rushing** through a pump opening cross-sectional area  **$5\text{cm}^2$**  at a speed of  **$12\text{m/s}$** , so the mass of the water coming out from the pump within **30 minutes** is .....

- ☐  **$18.2 \times 10^3 \text{ kg}$**
- ☐  **$15.1 \times 10^3 \text{ kg}$**
- ☐  **$10.8 \times 10^3 \text{ kg}$**
- ☐  **$8.6 \times 10^3 \text{ kg}$**




The **continuity equation** is represented  
by graph .....







In the opposite figure, the speed of the steady flow of water at A and C is **8 m/s** and **4 m/s** respectively, then its speed at B is .....



- ☐ 16 m/s
- ☐ 12 m/s
- ☐ 8 m/s
- ☐ 6 m/s

ہندذاکر  
آن لائن



**When the temperature of a liquid decrease. Its  
viscosity coefficient ...**

- ☐ Increases
- ☐ Decreases
- ☐ Does not change
- ☐ Cannot be determined without knowing the type of liquid





In the relatively low or medium speeds of a car, the air resistance due to its **viscosity** is .....

- ☐ Directly proportional to square the speed of the car
- ☐ Directly proportional to the speed of the car
- ☐ Inversely proportional to square the speed of the car
- ☐ Inversely proportional to the speed of the car.



**In the high speeds of a car, the air resistance due to its **viscosity** is .....**

- ☐ Directly proportional to the speed of the car
- ☐ Inversely proportional to the speed of the car
- ☐ Directly proportional to square the speed of the car
- ☐ Inversely proportional to square the speed of the car .



**Final exam**  
**Senior 2**



پروفیسر شریف

PHYSICS  
TEACHER




**Which of the following is affected in the light wave when it is diffracted?**

- ☐ Its frequency
- ☐ Its wavelength
- ☐ Its velocity
- ☐ Its propagation direction






In the opposite figure, a light ray falls from medium (a) at angle  $45^\circ$  on the separating surface with medium (b), where it deviates from its original path by an angle of  $45^\circ$ , so the relative refractive index between the two media ( ${}_b n_a$ ) equals



- ☐  $\sqrt{2}$
- ☐  $\frac{\sqrt{3}}{2}$
- ☐  $\frac{1}{\sqrt{2}}$
- ☐  $\frac{2}{\sqrt{3}}$



The opposite (displacement-time) graph represents a body that moves a simple harmonic motion, **so.....**



	The amplitude (cm)	The frequency (Hz)
a	0.1	4
b	0.05	2
c	0.1	0.5
d	0.05	0.25





A layer of a viscous liquid of thickness **3 cm** and viscosity coefficient  **$1.2 \text{ kg/m.s}$**  is confined between two horizontal parallel plates. If a tangential force of  **$1.6 \text{ N}$**  acts on the upper plate to move it with a uniform speed of  **$1 \text{ m/s}$** , then the area of the upper plate equals.....

☐  $0.02 \text{ m}^2$

☐  $0.03 \text{ m}^2$

☐  $0.04 \text{ m}^2$

☐  $0.05 \text{ m}^2$

ہندذاکر  
آن لائن



**We don't hear the sound of explosions that happen in the Sun, because .....**

- ☐ the location of the explosions is very far
- ☐ the sound propagates as transverse waves
- ☐ the sound propagates as electromagnetic waves
- ☐ the sound propagates as mechanical waves





If the angle of minimum deviation for a light ray that falls on one of the faces of an equilateral triangular prism is  $60^\circ$ , the refractive index of the prism material for the incident light equals.....

☐  $\sqrt{2}$

☐ 1.5


☐ 1.6

☐  $\sqrt{3}$

ہندذاکر  
آن لائن



The opposite figure shows four light rays that fall on an isosceles triangular prism of refractive index 1.5, so which of these rays changes its direction by  $180^\circ$ ?



☐ (1)

☐ (2)

☐ (3)

☐ (4)

ہندذاکر  
آن لائن





If **red** and **blue** light rays fall with the same angle of incidence  $\phi$  on the separating surface from the optically rarer medium to an optically denser medium, then the ratio **between** the angle of refraction of **red** light and the angle of refraction of **blue** light  $\left(\frac{\theta_r}{\theta_b}\right)$  in the optically denser medium is.....

- ☐ greater than 1
- ☐ less than 1
- ☐ equal to 1
- ☐ indeterminable



The cross-sectional areas of the two ends of a tube are  $0.005 \text{ m}^2$  and  $0.01 \text{ m}^2$ . If water flows through the tube steadily and the volume of the flowing water within 15 minutes is  $9 \text{ m}^3$ , then the speed of the water in.....



a



b



c



d


	The wide cross-section	The narrow cross-section
a	0.6 m/s	1.5 m/s
b	1 m/s	1.5 m/s
c	0.6 m/s	2 m/s
d	1 m/s	2 m/s





A plastic transparent plate of refractive index **1.5** is used to make an aquarium. If a light ray gets reflected from a fish inside the water and falls on the plastic plate at an angle of incidence  **$35^\circ$**  as in the opposite figure, therefore the **emergence angle** of the light ray to the air equals.....

[Knowing that:  $n_{\text{water}} = 1.33$ ]



☐  $30.57^\circ$

☐  $35.41^\circ$

☐  $49.72^\circ$

☐  $52.33^\circ$



In Young's double-slit experiment, a light ray falls on the double-slit where the distance between the two slits is **0.19 mm** and they are **90 cm** away from the observation screen. If the distance between the central fringe and the first bright fringe is  **$3 \times 10^{-3} \text{ m}$** , so the wavelength of the used light is.....

☐ 490 nm

☐ 520 nm

☐ 603 nm


☐ 633 nm

ہندذاکر  
اؤنلاین





The opposite figure represents the motion of a vibrating string, so the velocity of the string is **maximum at .....**



- ☐ point a
- ☐ point b
- ☐ points b and c
- ☐ points a and c



Four identical solid balls are **dropped from the same height** into four cylinders, each of them contains the **same volume** of **different liquid** while the time taken by each ball to reach the bottom of the cylinder is recorded as the following table:.....



Cylinder 1



Cylinder 2



Cylinder 3



Cylinder 4

Cylinder	Time
1	0.2 s
2	0.3 s
3	0.6 s
4	1 s





When a light of wavelength is used in Young's double-slit experiment, the path difference between the two interfered waves at the central fringe is equal to.....

☐  $1.5 \lambda$

☐  $\lambda$

☐  $0.5\lambda$

☐ 0

ہندذاکر  
آن لائن



A light ray falls on a mirror making an angle of  $60^\circ$  with its surface, therefore the reflection angle of the light ray from the mirror is equal to



$60^\circ$



$30^\circ$



$90^\circ$



$120^\circ$

ہندذاکر  
آن لائن





A blue light source is immersed under the surface of water at a certain depth to form blue light circular spot on the surface of water. If the blue light source is replaced by a red light source, then the light spot at the water surface .....

- ☐ vanishes
- ☐ decreases in area
- ☐ keeps its area
- ☐ increases in area



If the ratio between the apex angles of two thin prisms of the same material equals  $\frac{2}{5}$  then the ratio between the dispersive powers of them respectively equals .....

☐ 1 / 1

☐ 2 / 5

☐ 5 / 2

☐ 2 / 3

ہندذاکر  
آن لائن





If the volume flow rate of a liquid is  $Q_v$  in a pipe that is branched into four branches of equal **cross-sectional areas**, then the flow rate in each branch equals .....



$4 Q_v$



$1/3 Q_v$



$Q_v$




$1/4 Q_v$

ہندذاکر  
آن لائن



In the opposite figure, a tone of frequency **5000 Hz** is produced due to the vibration of a guitar string, then the periodic time of the vibrating string in ms equals .....



☐  $2 \times 10^{-4}$

☐  $5 \times 10^{-4}$

☐ 0.2

☐ 0.5

ہندذاکر  
آنونلااین





If a light ray falls on one of the faces of a triangular prism of apex angle  $40^\circ$  with an angle of incidence of  $60^\circ$  to emerge normally from the other face, then the refractive index of the prism equals

.....

- ☐ 1.5
- ☐ 1.41
- ☐ 1.35
- ☐ 0.71

ہندذاکر  
آنونلااین




People in the high floors feel wind speed more than those in the lower floors: **Explain why?**

هنذاكر  
أونلاين






The opposite graph shows the relation between the displacement ( $d$ ) and the time ( $t$ ) for two waves A and B, find the speed of propagation of each wave in the medium.





The opposite figure shows a light ray that falls on a triangular prism of refractive index **1.5**, trace the path of the light ray in the prism, then find the angle of emergence from the prism.





# كيفية طباعة صفحات معينة من ملف معين مثلا ازاي نطبع الصفحات من صفحة 4 الى صفحة 9



خطوة 1



خطوة 2  
اختيار اسم  
الطابعة  
بتاعتك

خطوة 3  
كتابة الصفحات  
المراد طباعتها  
نكتب رقم 4 ثم  
نكتب الشرطة  
دي - ثم نكتب 9

خطوة 4  
اختيار نوع الورق



خطوة 5  
اختيار A4



خطوة 6

حمل الآن

مجاناً وحصرياً

# المراجعة رقم (2)

## الترم الاول





## Important laws

1- **frequency** ( $\nu$ ) =  $\frac{n}{t} = \frac{1}{T} = \frac{1}{4 \times t_A} = \frac{v}{\lambda}$

2- **Periodic time** ( $T$ ) =  $\frac{t}{n} = \frac{1}{\nu} = 4 \times t_A = \frac{\lambda}{v}$

3- **Wavelength** ( $\lambda$ ) =  $\frac{\text{Total distance (X)}}{\text{Number of waves (n)}} = \frac{v}{\nu}$

4-  **$v = \lambda \cdot \nu$**

5-  $\frac{\lambda_1}{\lambda_2} = \frac{v_2}{v_1}$  &  $\frac{\lambda_1}{\lambda_2} = \frac{v_1}{v_2} = \frac{n_2}{n_1}$

6-  ${}_1n_2 = \frac{\sin \phi}{\sin \theta} = \frac{v_1}{v_2} = \frac{n_2}{n_1} = \frac{1}{{}_2n_1} = \frac{\lambda_1}{\lambda_2}$

7- **snell's law** ( $n_1 \sin \phi = n_2 \sin \theta$ )

8- The distance between two successive fringes of the same kind ( $\Delta y = \frac{\lambda R}{d}$ ), ( $\Delta y = \frac{x}{n}$ )

9-  $\sin \phi_c = \frac{n_2 (\text{less})}{n_1 (\text{high})} = {}_1n_2 = \frac{\sin \phi_{c1}}{\sin \phi_{c2}}$


When the less dense medium is air then:  $\sin \phi_c = \frac{1}{n}$

### Normal prism

10-  $A = \theta_1 + \phi_2$

11-  $\alpha = (\phi_1 + \theta_2) - A$

12-  $n = \frac{\sin \phi_1}{\sin \theta_1} = \frac{\sin \theta_2}{\sin \phi_2}$



At the minimum angle of deviation:

13-  $\phi_o = \frac{\alpha_o + A}{2}$  &  $\theta_o = \frac{A}{2}$

14-  $n = \frac{\sin(\frac{\alpha_o + A}{2})}{\sin(\frac{A}{2})}$

### The thin prism

15-  $\alpha_o = A (n - 1)$

16- **angular dispersion** =

$(\alpha_o)_b - (\alpha_o)_r = A (n_b - n_r)$

17-  $n_y = \frac{n_b + n_r}{2}$

18-  $(\alpha_o)_y = \frac{(\alpha_o)_b + (\alpha_o)_r}{2} = A (n_y - 1)$

\*If the thin prism immersed in a liquid then

$$\alpha_o = A \left( \frac{n_{\text{prism}}}{n_{\text{liquid}}} - 1 \right)$$

19-  $\omega_\alpha = \frac{(n_b - n_r)}{(n_y - 1)}$

## Chapter 4: Hydrodynamics

20-  $Q_v = \frac{v_{ol}}{t} = AV$

21-  $Q_m = \frac{\Delta m}{\Delta t} = \rho AV = \rho Q_v$

22-  $A_1 V_1 = A_2 V_2$

$$\therefore \frac{V_1}{V_2} = \frac{A_2}{A_1} = \frac{r_2^2}{r_1^2} = \frac{D_2^2}{D_1^2}$$

\*The tube is branched into (n) branches of the same area.

$$A_1 V_1 = n A_2 V_2 \quad \& \quad r_1^2 V_1 = n r_2^2 V_2$$

\* The tube is branched into (n) branches of different area.

$$* A_1 V_1 = A_2 V_2 + A_3 V_3 + A_4 V_4$$

$$* r_1^2 V_1 = r_2^2 V_2 + r_3^2 V_3 + r_4^2 V_4$$

### viscosity

23-  $F = \eta_{vs} \frac{AV}{d}$

24-  $\eta_{vs} = \frac{Fd}{AV}$

\*The measuring unit of viscosity coefficient:

N.s/m<sup>2</sup>, J.s/m<sup>3</sup>, Kg/m.s, Pascal.Sec.

### \*Conversions

Cm  $\longrightarrow$  m  $\times 10^{-2}$

mm  $\longrightarrow$  m  $\times 10^{-3}$

cm<sup>2</sup>  $\longrightarrow$  m<sup>2</sup>  $\times 10^{-4}$


cm<sup>3</sup>  $\longrightarrow$  m<sup>3</sup>  $\times 10^{-6}$

Angstrom =  $10^{-10}$  m



**12- A liquid flows steadily in a tube as in the opposite figure. If the cross-sectional areas of x, y, z are A, A, 2 A respectively and the liquid speeds at the cross-sections x, y are v, 2 v respectively, then the liquid speed at z equals.....**

- a)  $2v$                       b)  $\frac{2}{3}v$   
c)  $\frac{3}{2}v$                       d)  $v$



**13- The average speed of blood in one of the arteries of an adult person is 0.33 m/s. If the radius of that artery is 0.7 cm and it distributes blood to 30 smaller arteries each of radius 0.35 cm, then the average blood speed inside these small arteries equals.....**

- a) 0.011 m/s                      b) 0.022 m/s  
c) 0.033 m/s                      d) 0.044 m/s

**14- In steady flow, when the cross-sectional area of the tube decreases, the density of streamlines .....**

- a) increases                      b) decreases but doesn't reach zero  
c) vanishes                      d) remains unchanged

**15- The continuity equation of liquid flow can be deduced from the law of conservation of.....**

- a) mass                      b) energy  
c) momentum                      d) density

**16- If the cross-sectional area of a tube in which a liquid flows steadily gets increased to the double, then the volume flow rate.....**

- a) increases to the double                      b) decreases to its half  
c) remains constant                      d) decreases to its quarter

**17- Water flows steadily in a tube of diameter 2 cm at a speed of 5 m/s. Thus;**

**(i) The volume of water which is flowing through the cross-section of the tube in one minute equals.....**

- a)  $9.42 \text{ m}^3$                       b)  $0.19 \text{ m}^3$   
c)  $0.0942 \text{ m}^3$                       b)  $0.001 \text{ m}^3$

**(ii) The time required to fill a tank of volume  $20 \text{ m}^3$  by using the flowing water from the tube is.....**

- a) 127.38 minutes                      b) 212.31 minutes  
c) 3.54 minutes                      d) 2.123 minutes


**Mr. Abdullah Abdelazeem**



## Choose the correct answer

1- In the opposite figure: If the time taken by the pendulum to move from X to Z is 0.8 s, calculate:

- (a) The periodic time.
- (b) The frequency.
- (c) The number of complete oscillations through 16 s.
- (d) The time required to make 50 oscillations.



2- A sound wave of frequency 512 Hz travels from air to water. If the speed of sound in air is 340 m/s and in water is 1360 m/s, so the frequency of the wave in water equals.....

- a) 128 Hz
- b) 256 Hz
- c) 512 Hz
- d) 2048 Hz

3- If a vibrating object takes 0.1 s to complete one oscillation, then the number of complete oscillations that is made by the object during 100 s equals..... oscillations.


- a) 10
- b) 100
- c) 1000
- d) 10000

3- Sound waves travel in gases as.....

- a) longitudinal waves
- b) transverse waves
- c) longitudinal and transverse waves
- d) electromagnetic waves

4- In the opposite wave, which of the points a, b, c, d have the same phase?

- a) a,b,c
- b) a,b
- c) b,c
- d) b,d




5- If the time interval between the pass of the first crest and the tenth crest by a point in the path of a wave motion is 0.2s then the frequency is.....

- a) 45 Hz
- b) 50 Hz
- c) 55 Hz
- d) 60 Hz

6- The opposite graph shows the relation between the speed and the wavelength for two different waves of the same type (A and B) when they propagate through many different media, so.....

- a)  $v_A > v_B$
- b)  $v_A < v_B$
- c)  $\lambda_A = \lambda_B$
- d)  $\lambda_A > \lambda_B$



**Mr. Abdullah Abdelazeem**



7- If the speed of water waves that pass by a certain point is 1.5 m/s and there are 30 waves pass by this point in 1 s, then the number of waves in a distance of 60 m equals.....

- a) 3 waves                      b) 40 waves  
c) 400 waves                  d) 1200 waves

8- If the terminal of a spring is being moved to make a transverse wave of wavelength 30 cm and periodic time 0.1 s, then the terminal of the spring is being moved to make a longitudinal wave of periodic time 0.2 s which has the same speed as the transverse wave, so the wavelength of the longitudinal wave equals.....


- a) 7.5 cm                      b) 15 cm  
c) 30 cm                      d) 60 cm

9- A wave of frequency  $\nu_1$  and wavelength  $\lambda_1$ , propagates in a medium with speed  $v_1$ , if this wave travels from this medium to another medium where its speed becomes  $\frac{2}{3} v_1$ , then .....

- a) the frequency  $\nu_1$  remains constant and the wavelength becomes  $\frac{3}{2} \lambda_1$   
b) the frequency  $\nu_1$ , remains constant and the wavelength becomes  $\frac{2}{3} \lambda_1$   
c) the wavelength  $\lambda_1$  remains constant and the frequency becomes  $\frac{3}{2} \nu_1$   
d) the wavelength  $\lambda_1$  remains constant and the frequency becomes  $\frac{2}{3} \nu_1$


10- The opposite figure shows a wave which is travelling through medium x then it moves to another medium y, so the ratio between the speed of the wave in medium x to its speed in medium y ( $\frac{v_x}{v_y}$ ) is .....

- a) greater than one                      b) less than one  
c) one    d) indeterminable



11- The opposite figure shows a liquid flowing steadily in a tube from one terminal to the other, so the ratio of the numbers of streamlines through the cross-sections x : y : z is.....


- a) 2 : 1 : 4                      b) 1 : 2 : 4  
c) 2 : 4 : 1                      d) 1 : 1 : 1





**38- The opposite graph represents the variation of the distance between two bright consecutive fringes ( $\Delta y$ ) versus the reciprocal of the distance between the double slits ( $\frac{1}{d}$ ), if the distance between the double slit and the observation screen is 2 m, then the wavelength of the used light equals.....**

- a)  $5 \times 10^{-3}$  m                      b)  $2 \times 10^{-6}$  m  
c)  $10^{-6}$  m                          d)  $5 \times 10^{-7}$  m




**39- It is difficult to observe light diffraction in daily life because of.....**

- a) the high speed of visible light  
b) the small frequencies of visible light  
c) the short wavelengths of visible light  
d) the high intensity of visible light

**40- The opposite figure shows three transparent media(1) , (2) and (3), so if a light ray passes as shown in the figure, what will be the correct order of light speeds through these three media?**

- a)  $V_1 > V_2 > V_3$                       b)  $V_1 > V_3 > V_2$   
c)  $V_2 > V_3 > V_1$                       d)  $V_3 > V_2 > V_1$



**41- The critical angle between two media depends on.....**

- a) the absolute refractive index of the optically denser medium only  
b) the absolute refractive index of the optically rarer medium only  
c) the absolute refractive indices of the two media  
d) the angle of incidence of the light ray on the boundary surface between the two media


**42- The opposite figure represents a light ray that is incident at an angle of  $45^\circ$  on one of the faces of an equilateral triangular prism whose refractive index equals 1.5, so:**

**(i) The angle at which the ray emerges from the prism equals .....**

- a)  $28.13^\circ$                       b)  $31.87^\circ$   
c)  $42.12^\circ$                       d)  $52.37^\circ$

**(ii) The angle of deviation for the light ray equals.....**


- a)  $37.37^\circ$                       b)  $41.43^\circ$                       c)  $52.63^\circ$                       d)  $67.37^\circ$





30- The opposite figure shows a light ray falling on a glass cuboid to get refracted inside it, hence the speed of light ray in glass is approximately equal to..... (Given that:  $c = 3 \times 10^8$  m/s)

- a)  $1.8 \times 10^8$  m/s                      b)  $2 \times 10^8$  m/s  
a)  $2.3 \times 10^8$  m/s                      c)  $2.5 \times 10^8$  m/s



31- When a light ray falls on the interface between two media at an angle of incidence  $\phi$  and gets refracted at an angle of refraction  $\theta$ , the ratio  $\frac{\sin \phi}{\sin \theta}$  is.....

- a) constant for the two media                      b) variable according to the value of  $\phi$   
c) constant and always less than one                      d) constant and always greater than one

32- When the angle of incidence on the boundary surface between two media gets doubled, the relative refractive index between them.....

- a) decreases to its half                      b) gets doubled  
c) remains constant                      d) gets tripled

33- When a light ray is incident from air on water surface at an angle of incidence  $60^\circ$ , the angle of refraction will be.....

- a) greater than  $60^\circ$                       b) less than  $60^\circ$   
c) equal to  $60^\circ$                       d) equal to  $0^\circ$

34- A light ray is incident from air on the surface of a glass slab at an angle of  $52^\circ$ . If the ray is deviated by an angle of  $19^\circ$ , the refractive index of glass is.....

- a) 0.83                      b) 1.33  
c) 1.45                      d) 1.65

35- The double-slit in Young's double-slit experiment works as two coherent light sources, in which coherence means that the two waves formed from the double-slit have the same.....

- a) phase                      b) amplitude                      c) speed                      d) direction

36- In Young's double-slit experiment, the interference fringes widths increase when.....

- a) the distance between the double-slit and the observation screen decreases  
b) the distance between the double-slit and the observation screen increases  
c) the distance between the two slits increases  
d) the wavelength of the used monochromatic light decreases

37- In a Young's double-slit experiment, when increasing the intensity of the used light, the distance between the center of the central fringe and that of the first dark fringe.....

- a) increases                      b) decreases                      c) vanishes                      d) doesn't change



25- When a tangential force  $F$  acts on a plate of area  $A$  placed on another static plate where a layer of liquid of thickness  $d$  is in between, the upper plate moves with a uniform velocity  $v$ . What is the tangential force that makes the upper plate moves with a constant velocity  $2v$  under the same conditions?


- a)  $F$                       b)  $2F$   
c)  $\frac{F}{2}$                       d)  $\frac{F}{4}$

26- At relatively low or medium speeds of a car, the air resistance due to air viscosity is .....


- a) directly proportional to the square of the speed of the car  
b) directly proportional to the speed of the car  
c) inversely proportional to the square of the speed of the car  
d) inversely proportional to the speed of the car

27- A light ray is incident on a plane mirror with an angle of  $60^\circ$ . If the mirror is rotated by an angle of  $18^\circ$  in the clockwise direction while keeping the incident ray in the same direction as in the figure, then the angle of reflection becomes.....


- a)  $18^\circ$                       b)  $42^\circ$   
c)  $48^\circ$                       d)  $78^\circ$




28- The four following figures represent four rays of a light that has a wavelength  $\lambda$  travelling from medium A to other four different media x, y, z and k each one at a time. In which of these media (x, y, z or k) the light ray will have the longest wavelength?




(a)



(d)




(c)



(d)

29- The opposite figure represents with a definite drawing scale a red light ray while passing between two media, so the relative refractive index between those two media ( ${}_1n_2$ ) equals.....

- a) 1.44                      b) 1.18  
c) 1.13                      d) 0.88






18- Water flows steadily through a tube XY such that its speed and its mass flow rate at cross-section X are  $v$ ,  $Q_m$  respectively. If the water speed at cross-section y is  $\frac{v}{2}$ , then its mass flow rate is equal to.....

- a)  $2 Q_m$                       b)  $Q_m$   
c)  $\frac{Q_m}{2}$                          d)  $\frac{Q_m}{4}$

19- The graph that represents the continuity equation for liquids flow is.....



20- Oil flows in a tube x at a rate of 6 liters/minute and gets out from another tube y which is connected to the first tube x at a speed of 4 m/s, then the cross-sectional area of the second tube y equals.....

- a)  $1.5 \times 10^{-3} \text{ m}^2$                       b)  $1.5 \text{ m}^2$   
a)  $2.5 \times 10^{-5} \text{ m}^2$                       b)  $0.025 \text{ m}^2$

21- A plane surface of area  $0.5 \text{ m}^2$  moves at a uniform velocity of 2 m/s parallel to another static surface that is separated from it by a layer of liquid of thickness 4 cm, if the viscosity coefficient of the liquid is  $1.5 \text{ kg/m.s}$ , then the force required to keep the moving with this uniform velocity is.....

- a) 37.5 N                      b) 50 N  
c) 67.5 N                      d) 150 N



22- If the speed difference between two liquid layers gets decreased when a tangential force is acting on the upper layer, then at the same temperature the viscosity coefficient.....

- a) vanishes                      b) decreases but doesn't vanish                      c) increases                      d) remains constant

23- When the temperature of a liquid decreases, its viscosity coefficient.....

- a) increases                      b) decreases  
c) does not change                      d) depends on the type of liquid

24- Which of the following graphs represents the relation between the force (F) which is required to move plate x with a uniform velocity  $v$  on the surface of a liquid parallel to a static plate y and  $t$




**Mr. Abdullah Abdelazeem**



43- The opposite figure shows a light ray which is incident normally on one of the faces of a triangular prism of an apex angle  $40^\circ$ , if  $\theta_2 = 1.5\theta_1$  the refractive index of the prism equals.....

- a) 1.25                      b) 1.35  
c) 1.51                      d) 1.72




44- In the opposite figure, a blue light ray falls on the face of a prism at point P so that the angle of refraction equals  $23^\circ$  then it falls on the opposite face at point Q and emerges tangent to that face, hence:

(i) The critical angle of the prism's material for the blue light equals.....

- a)  $23^\circ$                       b)  $37^\circ$                       c)  $42^\circ$                       d)  $60^\circ$


(ii) The refractive index of the prism's material for the blue light equals

- a) 1.15                      b) 1.41  
c) 1.66                      d) 1.72



45- In the opposite figure, the apex angle (A) of the prism is.....

- a) greater than  $45^\circ$                       b) less than  $45^\circ$   
c) equal to  $45^\circ$                       d) indeterminable



46- A light ray falls normally on one of the faces of a triangular prism of apex angle  $30^\circ$ . If the refractive index of the prism is  $\sqrt{2}$  then:

(i) The angle of emergence of the ray from the prism equals.....

- a)  $15^\circ$                       b)  $30^\circ$   
c)  $45^\circ$                       d)  $60^\circ$

(ii) The angle of deviation of the ray equals .....

- a)  $15^\circ$                       b)  $30^\circ$   
c)  $45^\circ$                       d)  $60^\circ$

47- Two thin prisms x and y are made of the same material, the apex angle of prism x is A and its dispersive power is  $\omega_\alpha$  so if the apex angle of prism y is  $1.5A$ , its dispersive power is.....

- a)  $\frac{\omega_\alpha}{2}$                       b)  $\omega_\alpha$                       c)  $2\omega_\alpha$                       d)  $3\omega_\alpha$


**Mr. Abdullah Abdelazeem**

## Answer the following

1- The opposite figure shows a vibrating string that takes 0.01 s to reach a maximum displacement away from its rest position. Calculate the amplitude and the frequency of the string.


2- The opposite graph represents a transverse wave, calculate:

- (a) The amplitude.
- (b) The frequency.
- (c) The periodic time.
- (d) The wavelength.
- (e) the wave speed



3- The opposite graph represents a transverse wave, calculate:

- (a) The periodic time.
- (b) The frequency
- (c) The wavelength.
- (d) the wave speed



4- If a sound wave travelled from one medium to another and the ratio between its wavelengths in the two media was  $\frac{2}{3}$  respectively, what is the ratio between the speeds of sound in the two media respectively?



5- If the ratio between the frequency of a man's sound and that of a girl's sound is  $\frac{3}{4}$ , what is the ratio between the speeds of the sounds of the man and the girl in air?

6- A water pipe of diameter 2 cm in which water flows with a speed of 0.1 m/s. The pipe enters a house, where its diameter becomes 1 cm. Given that the density of water is  $1000 \text{ kg/m}^3$ , calculate:

(a) The speed of water in the pipe inside the house.


(b) The volume and the mass of the water that flows every minute through any cross-section of the pipe.

7- If the absolute refractive index of water is  $\frac{4}{3}$  and the absolute refractive index of glass is  $\frac{3}{2}$  calculate:

(a) The relative refractive index from water to glass.

(b) The relative refractive index from glass to water.

8- The opposite figure shows three mirrors A, B and C. If a light ray falls on mirror A as shown in the figure, trace the path of the ray till it gets reflected from mirror C.



9- A light ray falls on the surface of a glass of refractive index 1.5 at an angle of incidence  $60^\circ$ , if a small part of the light gets reflected and another part gets refracted, calculate the angle between the reflected and the refracted light rays.

10- In the double-slit experiment; if the distance between the two narrow rectangular slits was 0.15 mm, the distance between the double-slit barrier and the observation screen was 75 cm and the distance between the centers of two successive bright fringes was 0.3 cm, calculate the wavelength of the used monochromatic light source.

**36- A thin prism has an apex angle of  $8^\circ$ , its refractive index for red light is 1.52 and its refractive index for blue light is 1.54. Calculate:**

- (a) The angle of deviation for each light color.
- (b) The angular dispersion for the light in the prism.
- (c) The dispersive power of the prism.

## Exercise

**1- In the opposite figure: If the pendulum takes 1 s to move from point X to point Y, so its frequency equals.....**

- a) 0.5 Hz
- b) 5 Hz
- c) 10 Hz
- d) 50 Hz

**2- In the opposite figure, a flexible oscillating rod takes time of 0.01 s to move from a to b, then :**

**(i) Its periodic time.....**

- a) 0.02 s
- b) 0.04 s
- c) 0.06 s
- d) 0.08 s

**(ii) Its frequency is.....**



- a) 1.25 Hz
- b) 18.75 Hz
- c) 25 Hz
- d) 31.25 Hz

**(iii) The amplitude of its oscillation is.....**

- a) 3 cm
- b) 6 cm
- c) 9 cm
- d) 12 cm


**3- In the opposite figure, a liquid flows steadily in a tube, so the physical quantity which is greater at cross-section x than at cross-section y is.....**

- a) the liquid speed
- b) the volume of the flowing liquid in unit time
- c) the mass of the flowing liquid in unit time
- d) the number of the streamlines through the cross-section





17- In the opposite figure, if the refractive index of the prism is 1.5, calculate the deviation angle of the light ray.




18- A triangular prism is made of a material whose refractive index is  $\sqrt{2}$  having an apex angle that equals  $60^\circ$ , calculate:

- The minimum angle of deviation in the prism.
- The angle of incidence and the angle of emergence at minimum deviation.

19- The opposite graph represents the relation between the angle of deviation ( $\alpha$ ) for a light ray passing through a triangular prism and the first angle of incidence ( $\theta_1$ ) on the face of the prism, calculate:

- The apex angle of the prism.
- The refractive index of the prism.
- The angle of emergence from the prism at point x.




20- A thin prism has an apex angle of  $7^\circ$  and refractive index of 1.5. Calculate the angle of deviation of light in the prism.

21- When a red light ray fell on a thin prism of apex angle  $A$  and refractive index  $n$ , the ray deviated by an angle of  $4^\circ$ . If the prism has been submerged in a liquid of refractive index 1.2, the angle of deviation becomes  $2^\circ$ , Calculate:

- The absolute refractive index of the prism ( $n$ ).
- The apex angle of the prism ( $A$ ).

11- The opposite figure represents the interference pattern of Young's experiment which was conducted with a light of wavelength  $5000 \text{ \AA}$  and an observation screen at distance  $120 \text{ cm}$  from the double-slit. If the distance between the central fringe (0) and the fourth bright fringe (4) was  $0.8 \text{ cm}$ , calculate the distance between the two slits.



12- If the absolute refractive indices of glass and water for a given monochromatic light ray are  $1.6$  and  $1.33$  respectively, calculate:

- The critical angle for each of them with air.
- The critical angle for the incident light ray that travels from glass to water.


13- The speeds of propagation of a light wave through two different media (x and y) are  $2 \times 10^8 \text{ m/s}$  and  $2.75 \times 10^8 \text{ m/s}$  respectively. Calculate the critical angle between the two media.

14- A piece of diamond was placed at the bottom of a wide basin filled with water for a height of  $1 \text{ m}$ , calculate the smallest diameter of a cork disk that while floating on the water surface will be enough to block the reflected light by the diamond from emerging out of the water surface. (Giving that: The absolute refractive index of the water =  $1.33$ )

15- A light ray fell from glass on the boundary surface with water, so its wavelength changed from  $5000 \text{ \AA}$  to  $5625 \text{ \AA}$ , calculate the critical angle from glass to water.

16- In the opposite figure, if the refractive index of the prism's material is  $1.5$ :

- Trace the light ray inside the prism.
- Find the angle of emergence from the prism.
- Find the angle of deviation.



**Mr. Abdullah Abdelazeem**



29- In the minimum deviation position in prism the sum of apex angle and deviation angle equal.....

- a) Double the refraction angle
- b) Double the incidence angle
- c) Double the emergence angle
- d) b and c

30- The mirage phenomenon is from application of

- a) density
- b) viscosity
- c) the total internal reflection
- d) steady flow

31- A light ray fall from air on a triangular prism by angle 65 and emerge normal to the other side then the apex angle .....65


- a) greater than
- b) equal
- c) less than

32- Oscillating body its frequency is 25Hz so the time of amplitude is ..... sec.

- a) 0.01
- b) 0.02
- c) 0.04
- d) 0.06


33- The opposite figure show the transfer of light between X and Y then.....

- a)  $n_y = \frac{\sin\theta}{\sin\phi}$
- b) The speed of light in X is greater than Y
- c) The speed of light in X is less than Y
- d) The medium X has greater optical density than Y



34- In the opposite figure a light ray fall perpendicular and emerge by angle 40° then its deviation angle ..... 40°

- a) greater than
- b) equal
- c) less than



35- The time taken by a pendulum to pass by two points where its speed at one of them is maximum and zero at the other is 0.1s so its frequency equal.....

- a) 2.5 Hz
- b) 0.05 Hz
- c) 30 Hz
- d) 20 Hz

36- A thin prism its apex angle is 6° make a deviation equal 3.6° then its refractive angle is .....

- a) 1.5
- b) 1.6
- c) 1.7
- d) 1.8

**Mr. Abdullah Abdelazeem**

**4- The opposite graph represents the relation between the vertical displacement of one of the medium particles (d) and the time (t) for a wave, then:**

**(i) The amplitude of that wave is.....**

- a) 9 cm                      b) 17.5 cm
- c) 18 cm                    d) 35 cm

**(ii) The frequency of the wave is.....**


- a) 1.7 Hz                    b) 2.5 Hz
- c) 3.3 Hz                    d) 5 Hz

**(iii) The wavelength of the wave is.....**





- a) 0.05 cm                  b) 17.5 cm
- c) 20 cm                    d) 35 cm

**(iv) The propagation speed of the wave is.....**

- a) 0.5 m/s                    b) 0.6 m/s
- c) 50 m/s                    d) 60 m/s



**5- A transverse wave is propagating in different media and the following graphs represent the relation between the vertical displacement (d) of the medium particles at a certain moment and the propagation distance (x) of the wave with the same scale. In which medium does the wave have the highest speed?**





6- A stone was thrown into a lake, so 50 waves were formed after 5 seconds from the collision of the stone with the water, when the radius of the outer circle was 2 m, then:

(i) The wavelength of the wave is.....

- a) 0.04 m                      b) 0.08 m
- c) 25 m                        d) 100 m

(ii) The frequency of the wave is.....

- a) 0.1 Hz                      b) 10 Hz
- c) 25 Hz                      d) 250 Hz

(iii) The propagation speed of the wave is.....

- a) 2.5 m/s                      b) 2 m/s
- c) 1 m/s                        d) 0.4 m/s

7- A liquid of density  $1000 \text{ kg/m}^3$  flows steadily inside a tube of cross-sectional area  $0.5 \text{ m}^2$  with a flow rate of  $10 \text{ kg/s}$ , then the flow speed of the liquid through the tube is equal to.....

- a) 200 m/s                      b) 50 m/s
- c) 0.02 m/s                      d) 0.05 m/s

8- Water flows steadily in a tube of cross-sectional area  $10^{-3} \text{ m}^2$ , if the volume of water which is coming out from the tube within 30 minutes is  $18 \text{ m}^3$  then.....


	Volume flow rate ( $\text{m}^3/\text{s}$ )	Speed of water flow ( $\text{m/s}$ )
(a)	0.01	10
(b)	0.01	600
(c)	0.6	10
(d)	0.6	600

9- If the ratio between the radii of two cross-sections of a tube in steady flow is  $\frac{1}{2}$ , then the ratio between the speeds of the liquid through them respectively is.....

- a)  $\frac{1}{4}$                       b)  $\frac{1}{2}$
- c)  $\frac{2}{1}$                       d)  $\frac{4}{1}$

10- In the opposite figure, the tubes (A) and (B) are different in cross-sectional area and the steady flow rate of the liquid inside each of them is  $0.3 \text{ m}^3/\text{s}$ . The two tubes meet to open in the tube (C) as in the figure, then the volume flow rate in the tube (C) is.....

- a)  $0.1 \text{ m}^3/\text{s}$                       b)  $0.3 \text{ m}^3/\text{s}$
- c)  $0.6 \text{ m}^3/\text{s}$                       d)  $0.9 \text{ m}^3/\text{s}$



21- If the refractive index of medium X is double that of Y, so the speed of light in X .....that in Y.


- a) Double                      b) half                      c) quarter                      d) 3 times

22- If the speed of wave is changed due to the change of medium this mean.....


- a) Its frequency increase                      b) its periodic time increase  
c) its wavelength doesn't change                      d) its periodic time doesn't change

23- The opposite figure show the change of potential energy with time of a pendulum started its motion from .....


- a) Maximum displacement                      b) Maximum speed  
c) Maximum kinetic energy                      d) minimum potential energy




24- the best graph show the relation between the speed of light in air when its frequency change is .....




(a)



(b)



(c)



(d)

25- Two sound wave propagate in air their frequencies is 512Hz , 256Hz so the ratio between their velocities is.....

- a) 1:2                      b) 2:1                      c) 1:3                      d) 1:1

26- The ratio between the angle of incidence in glass ( $n_g = 1.5$ ) to the angle of refraction in water ( $n_w = 1.3$ ).....

- a) less than one                      b) greater than one                      c) equal one

27- When light travel from high dense medium to less dense medium, so the maximum angle of refraction in the less medium is.....

- a)  $180^\circ$                       b)  $42^\circ$   
c)  $45^\circ$                       d)  $90^\circ$

28- The ratio between the refractive index of red light to that of the violet light .....one.

- a) greater than                      b) equal                      c) less than




**17- The opposite figure :**

**Illustrates the relation between the displacements and the time for a wave .**

**Calculate:**

- a) The wavelength
- b) The wave velocity
- c) Periodic time
- d) Amplitude

(0.4444m ,  $11.11 \times 10^3 \text{ m/s}$  ,  $4 \times 10^{-5} \text{ S}$  , 4cm)




**18- A stone was thrown in a lake so 50 waves were formed after 5 second from the colliding of the stone with the water where the radius of the outer circle is 2m find:**

- a) The wavelength
- b) The frequency
- c) The wave velocity
- d) The periodic time


(0.04 m, 10 Hz, 0.4 m/s, 0.1 S)

**19- The opposite figure show the change of displacement of an oscillating object then the best graph that express the change in velocity of this object is.....**



**20- In the opposite figure the frequency is 8Hz, so the speed of this wave in m/s is.....**

- a) 0.64
- b) 0.32
- c) 6.4
- d) 3.2



**Mr. Abdullah Abdelazeem**

11- Blood flows through an artery with an average speed of 0.24 m/s. If the artery branches into 120 smaller arteries each of diameter that is  $\frac{1}{4}$  of the big one, then the flow speed of blood in every small artery equals.....

- a)  $8 \times 10^{-3}$  m/s      b) 0.08 m/s  
c) 0.032 m/s      d) 0.3 m/s


12- A layer of thickness  $x$  of a liquid of viscosity 0.2 kg/m.s is confined between two plates, one of the plates is static and the other which has an area of  $2 \text{ cm}^2$  is moving with a uniform velocity such that it covers a distance  $100x$  through a time interval of 4 s, so the required force to move the plate equals.....

- a) 10 N      b)  $10^{-3}$  N  
c)  $10^{-4}$  N      d) 0.1 N

13- A car was travelling with high speed on a desert road when its driver noticed that the fuel was about to run out, what is the best strategy to save the fuel until reaching the nearest fuel station, considering what you have studied?

14- The opposite figure shows two metal balls (A, B) falling into two identical cylinders containing equal volumes of the same liquid with different temperatures, which of the two balls reaches the bottom of the cylinder first?

- a) Ball A      b) Ball B  
c) The two balls reach the bottom at the same instant  
d) The two balls will never reach the bottom




15- A patient is injected by a needle of radius 0.3 mm, if the rate of the drug flow in cross-section of the needle is  $0.5 \text{ cm}^3/\text{s}$ , then the speed of the drug flow in the needle is .....

- a) 1.24 m/s      b) 1.77 m/s  
c) 2.42 m/s      d) 7.71 m/s

16- The opposite figure shows a load that is attached to a vibrating spring, so the amplitude of the vibration equals.....

- a) 3 cm      b) 6 cm  
c) 9 cm      d) 12 cm





**51- In one prism:**

- a) Its apex angle change according to the wavelength of light fall on it
- b) Has a constant refractive index whatever the wavelength of light change
- c) Its refractive index change by change the wavelength of light falling on it
- d) The refractive index change by change the angle of incidence of light on it


**52- In the opposite figure the medium which has maximum optical density.....**

- a) A
- b) B
- c) C
- d) D




**53- The opposite figure represents equilateral prism, so its refractive index is.....**

- a)  $\frac{1}{2}$
- b)  $\sqrt{2}$
- c)  $\sqrt{3}$
- d)  $\frac{\sqrt{3}}{2}$



**54- If a light ray fall as in the opposite figure then its apex angle.....**

- a) greater than the critical angle
- b) Smaller than the critical angle
- c) Equal the critical angle
- d) There is no relation with the critical angle



**55- The ratio between the mass flow rate to volume flow rate for the same liquid equal.....**

- a) density of liquid
- b) viscosity coefficient
- c) speed of liquid

37- when the apex angle of a thin prism increase so its dispersive power .....

- a) increase                      b) decrease                      c) remain constant

38- In Young's experiment the path difference at the central fringe is.....

- a) 0                      b)  $2\lambda$                       c)  $\lambda$                       d)  $0.5\lambda$

39- The ratio between the wavelength of light in air to the wavelength of same light in water .....one.

- a) greater than                      b) equal                      c) less than


40- If the distance from the first crest to the third trough in a transverse wave = 25cm ,so the wavelength =.....cm.

- a) 12.5                      b) 10                      c) 15                      d) 5

41- When using red light instead of the green light, so the number of fringes in a unit length that is formed on the observation screen.....

- a) Increase                      b) decrease                      c) increase then decrease                      d) doesn't depend on color


42- When red and blue light rays on a thin prism show which of the following relations is perfect to express the relation between the deviation angle and the apex angle.



43- When replacing source of red light under water surface by other blue, so the radius of the light appear on the water surface.

- a) increase                      b) decrease                      c) doesn't change                      d) can't determine it

44- When replace a thin prism (A) by other (B) its apex angle is smaller than that of (A), so the right graph between the deviation angle and the refractive index is.....



**Mr. Abdullah Abdelazeem**



45- In the optical fiber the optical density of the outer layer ..... the optical density of the inner layer

- a) greater than                      b) equal                      c) less than

46- The incident light rays differ than the reflected rays in.....

- a) frequency                      b) speed                      c) wavelength                      d) intensity


47- In the opposite graph :

(i) The result of dividing (a) on (b) = .....


- a)  $\frac{-1}{A}$                       b)  $-A$                       c)  $-1$                       d)  $A$

(ii) The result of dividing the value of slope on value of point (a)

- a)  $\frac{-1}{A}$                       b)  $-A$                       c)  $-1$                       d)  $A$




48- Three different mediums a light ray fall on it choose the medium which has greater critical angle.



49- Find the wavelength of light of frequency  $5 \times 10^{14}$  Hz when propagate in diamond of refractive index  $\frac{5}{2}$  (knowing that the speed of light in space is  $3 \times 10^8$  m/s) (240  $\times 10^{-9}$  m)

50- In the opposite figure choose the correct answer

- a) the optical density of (1) is greater than (2)  
b) angle of incidence is less than the angle of refraction  
c) the light ray may emerge tangent if it is fall from medium (1) to medium (2)  
d) the light ray may emerge tangent if it is fall from medium (2) to medium (1)



حمل الآن

مجاناً وحصرياً


# امتحانات رقم (1)

## الترم الاول






1. A light ray fell from medium (1) its absolute refractive index is 1.3 to medium (2) its absolute refractive index is 1.5 as shown in the figure.



Which one from the following choices shows what happens for both wavelength and the velocity of the light wave in medium (2)?

	Wavelength	Velocity
A	Increases	Increases
B	Decreases	Increases
C	Increases	Decreases
D	Decreases	Decreases

2. The graph represents a wave motion with wavelength ( $\lambda$ )




The horizontal distance (x) between the two points A and B represents

.....

- A.  $\frac{3\lambda}{2}$   
B.  $\frac{2\lambda}{3}$   
C.  $2\lambda$   
D.  $\lambda$

3. The graph represents the relation between the displacement of one from medium particles during a certain time (d) and the covered distance by the wave during the same time (x).



Which one from the following choices represents the wave amplitude and wavelength?


	Wave amplitude	Wavelength
A	The distance KL	The distance mn
B	A half of distance KL	A double of distance mn
C	A double of distance KL	The distance mn
D	A half of distance KL	A half of distance mn



4. In Young's experiment to study the light interference, a monochromatic ray its wavelength  $6000 \text{ \AA}$  was used forming fringes on an observing screen at distance (R) from the double slits and the distance between two successive bright fringes was ( $\Delta y_1$ ). If another monochromatic light its wavelength  $4000 \text{ \AA}$  is used and the distance between the double slits and the observing screen is doubled, the distance between two successive bright fringes becomes ( $\Delta y_2$ ), the ratio  $\frac{\Delta y_1}{\Delta y_2} = \dots\dots\dots$

- A.  $\frac{3}{4}$
- B.  $\frac{4}{3}$
- C.  $\frac{6}{4}$
- D.  $\frac{1}{3}$

5. Five light rays are separated by equal angles each angle is  $15^\circ$  as shown in the figure if these light rays fall from medium (x) its absolute refractive index is 1.5 to medium (Y) its absolute refractive index is 1.33



**How many rays can penetrate to medium (Y)?**

- A. Four rays
- B. Three rays
- C. Two rays
- D. Five rays

6. Two thin prisms made of same material their apex angles ( $5^\circ$  and  $10^\circ$ ),

**the ratio between their dispersive power**  $\frac{(w_{\alpha 0})_1}{(w_{\alpha 0})_2} = \dots\dots\dots$

- A. 0.5
- B. 0.6
- C. 1
- D. 2

7. A light ray falls with angle of ( $45^\circ$ ) on one side of a triangular prism its apex angle is  $30^\circ$  and emerged normally from the opposite side, **so the angle of deviation** = .....

- A.  $15^\circ$
- B.  $20^\circ$
- C.  $25^\circ$
- D.  $30^\circ$



8. Four identical metallic balls are fallen down from the same height, each ball toward a separate jar where each jar contains a different liquid from the other jars then the taken time by each ball to reach the bottom of the jar is recorded in each case as the following:

The jar	The time taken by the ball to reach the bottom
1	0.2 s
2	0.3 s
3	0.6 s
4	1 s

**Which jar contains the liquid with higher viscosity?**

- A. Jar 1
- B. Jar 2
- C. Jar 3
- D. Jar 4

9. A liquid flows with velocity ( $v$ ) through a uniform tube, its diameter is ( $X$ ) and a cork has small aperture is put at one end of this tube. If the diameter of the small aperture is ( $\frac{X}{4}$ ). **The velocity of emergent water from the small aperture = .....**

- A.  $16 v$
- B.  $4 v$
- C.  $\frac{1}{4} v$
- D.  $\frac{1}{16} v$

10. A wave its frequency (100 Hz) propagates in air with velocity of 320m/s, **calculate its wavelength.**

.....

.....

.....

11. A light wave propagates with velocity  $2 \times 10^8$  m/s through a certain medium its refractive index 1.5 if this light wave transfers to another medium its refractive index is 1.2, **Find the velocity of light in the second medium.**

.....

.....

.....

12. A thin prism of apex angle  $10^\circ$ , its absolute refractive index of both blue and red colors (1.53 and 1.51) respectively.  
**Calculate the average deviation angle.**

.....

.....

.....



# كيفية طباعة صفحات معينة من ملف معين مثلا ازاي نطبع الصفحات من صفحة 4 الى صفحة 9



خطوة 1



خطوة 2  
اختيار اسم  
الطابعة  
بتاعتك

خطوة 3  
كتابة الصفحات  
المراد طباعتها  
نكتب رقم 4 ثم  
نكتب الشرطة  
دي - ثم نكتب 9

خطوة 4  
اختيار نوع الورق



خطوة 5  
اختيار A4



خطوة 6